



Rx FOR SUCCESS

Transient Ischemic Attacks (TIAs) and Strokes (CVAs)

Obstruction in blood flow (ischemia) to the brain can lead to permanent damage. This is called a cerebrovascular accident (CVA). It is also known as cerebral infarction or stroke. If the symptoms are temporary without permanent brain damage, the event is called a transient ischemic attack (TIA). Rupture of an artery with bleeding into the brain (hemorrhage) is called a CVA, too. Strokes and TIAs are rated based on the underlying cause.

The most common cause of TIAs and CVAs is hypertensive and atherosclerotic plaque within the arteries to the brain (aka cerebrovascular disease or CVD). CVD can be complicated by clots (thrombosis) and by emboli from the heart. Because CVD is an indicator of atherosclerosis in other parts of the body, an individual with a history of TIA or CVA is at risk for coronary artery disease and recurrent stroke. Risk factors for CVD include smoking, coronary artery disease, high blood pressure, diabetes, lipid disorders (such as high cholesterol), peripheral arterial disease, and atrial fibrillation.

Signs and symptoms of a CVA/TIA include weakness, numbness, headaches, dizziness, nausea, vomiting, paralysis of one side of the body, speech difficulty, and memory defects. Amaurosis fugax, a form of visual TIA, is temporary monocular (one eye) or partial blindness.

Tests are done to evaluate the brain circulation, such as a carotid ultrasound (Duplex) or angiogram (MRA). A brain scan (CT and/or MRI) is used to determine if an individual has had a stroke. A TIA will not show on a scan. TIA is never ruled-out by negative tests; diagnosis is adequately met by symptoms only.

Although CVAs, TIAs, and bleeding into the brain are mainly due to atherosclerosis or hypertension, there are many non-atherosclerotic causes: migraine, adverse drug reactions, trauma, ruptured congenital aneurysm, valvular heart disease, congenital heart disease, clotting disorders, connective tissue disease (example lupus), and others. At times, no cause can be found in young individuals. This is termed "cryptogenic." Cryptogenic events over age 55 are assumed to be atherosclerotic. The long term prognosis varies depending on the cause, and additional tests (such as echocardiogram, clotting studies, and other blood and imaging tests) may be required to determine rare etiologies.

Treatment for CVD includes physical and speech rehabilitation for any residual impairment, blood thinners (like aspirin or Coumadin), cholesterol lowering medications, and blood pressure control. Surgical treatment (endarterectomy or stent) may be used to open the obstruction.

This material is designed to provide general information about the subject matter covered. It should be used with the understanding that we are not rendering legal, accounting, or tax advice. Such services should be provided by the client's own professional advisors. Accordingly, any information in this document cannot be used by any taxpayer for purposes of avoiding penalties under the Internal Revenue Code.

This material is intended for insurance informational purposes only and is not personal medical advice for clients. Rates and availability will vary based on the satisfaction of our underwriting criteria. Underwriting rules are subject to change at our discretion. This marketing material is subject to an expiration date, and use of this material must be discontinued as of the expiration date.

Insurance issued by The Prudential Insurance Company of America and its affiliates, Newark, NJ.

NOT FOR CONSUMER USE.

© 2013 Prudential Financial, Inc. and its related entities.
0191521-00003-00 Ed. 02/2013 Exp. 02/13/2015 Rx 016

Securities and Insurance Products:

Not Insured by FDIC or Any Federal Government Agency.
May Lose Value.

Not a Deposit of or Guaranteed by Any Bank or Bank Affiliate.



Prudential
Bring Your Challenges[®]

A general idea of ratings for TIA/CVA can be gotten from the tables below. Adjustments up or down in ratings will depend on the underlying cause, the extent of permanent neurological impairment, treatment, and the quality of risk factor modification. Acceptable cases have minimal residuals, normal mentation, and a return to full activity. More than two events, dementia, or significant residual physical or mental impairment are declined.

SINGLE EVENT DUE TO ATHEROSCLEROTIC DISEASE*			
Age at application	Younger than 45 years old	45 to 54 years old	55 years and older
TIA more than 6 months from event	Table B	Table B	Table B
CVA more than 6 months from event	Decline	Table D	Table C
TIA or CVA less than or equal to 6 months from event	Postpone	Postpone	Postpone
SINGLE EVENT PLUS ONE ADDITIONAL EVENT Add single event rating above to appropriate rating below.			
Age at application	Younger than 45 years old	45 to 54 years old	55 years and older
TIA more than 1 year from event	Non-Smoker Plus	Non-Smoker Plus	Non-Smoker Plus
CVA more than 1 year from event	Decline	Table C	Table C
CVA, significant residuals but still physically independent with normal mentation	Decline	An additional 1-2 tables	An additional 1-2 tables
*Credits of 1-2 tables are available for at least 2 years of stability after evaluation, treatment, and recovery.			

SINGLE EMBOLIC EVENT DUE TO HEART DISEASE	
Valve disease	Rate for valve disease, no less than Table B (See <i>Rx for Success</i> on Aortic Valve Disorders and Mitral Valve Disorders)
Atrial fibrillation	Sum AF and TIA/CVA debits (See <i>Rx for Success</i> on Atrial Fibrillation)

OTHER SINGLE EVENTS, FULLY RECOVERED	
TIA/CVA due to Migraine	Individual Consideration Table B or better
Cryptogenic event, trauma, adverse drug reaction (no longer on the offending drug)	Postpone one year, then 0
Clotting disorder, congenital heart malformation, others	Table ratings vary

Non-atherosclerotic causes of stroke include aneurysms, vascular malformations, trauma, clotting disorders, emboli from abnormal heart structures, vasculitis/arteritis, adverse drug events (warfarin, birth control pills, cocaine, amphetamine, etc.), fibromuscular dysplasia, spontaneous dissection. Risk assessment in Life underwriting depends on the underlying cause.

To get an idea of how a client with TIA or CVA would be viewed in the underwriting process, use the attached Ask "Rx"pert Underwriter for an informal quote. We also recommend Rx for Success on related topics: *Vascular Lesions of the Brain and Carotid Ultrasound and Brain Scan*.

Ask "Rx"pert Underwriter (Ask Our Expert)

After reading the *Rx for Success* on TIAs and CVAs, use this form to Ask "Rx"pert Underwriter for an informal quote.

Producer _____ Phone _____ Fax _____
 Client _____ Age/DOB _____ Sex _____

If your client has had TIA or CVA, please answer the following:

1. Please provide date(s) of each event.

2. What was the underlying cause of the TIA/CVA?

- Atherosclerosis and/or hypertension
- Atrial fibrillation
- Heart valve disease
- Congenital heart malformation, such as a hole in the heart
- Other. Please specify. _____

3. Is there a history of any other cardiovascular disease (e.g., peripheral arterial disease, carotid artery disease, coronary artery disease, etc.)?

- Yes. Please give details. _____
- No

4. Have any tests been completed? (Check all that apply.)

- Echocardiogram (Date) _____
- Carotid ultrasound/Duplex (Date) _____
- Brain scan by CT and/or MRI (Date) _____

5. Is your client on any medications?

- Yes. Please give details. _____
- No

6. Has your client smoked cigarettes in the last 12 months?

- Yes
- No

7. Does your client have any other major health problems (e.g., cancer, heart disease, seizures, psychiatric illness, etc.)?

- Yes. Please give details. _____
- No